

# INVERSION OF ADJUNCTION ON LOG CANONICITY

MASAYUKI KAWAKITA

ABSTRACT. We prove inversion of adjunction on log canonicity.

We work over an algebraically closed field of characteristic zero. Let  $X$  be a normal variety and  $B$  an effective  $\mathbb{R}$ -divisor on  $X$ . We call the pair  $(X, B)$  a *log pair* if  $K_X + B$  is an  $\mathbb{R}$ -Cartier  $\mathbb{R}$ -divisor. A valuation  $v$  of the function field of  $X$  is called an *algebraic valuation* of  $X$  if it is defined by a prime divisor  $E$  on a non-singular variety  $\bar{X}$  equipped with a birational morphism  $f: \bar{X} \rightarrow X$ . Such a valuation is denoted by  $v_E$ . If  $(X, B)$  is a log pair, we can write  $K_{\bar{X}} + \bar{B} = f^*(K_X + B) + A$ , where  $\bar{B}$  is the birational transform of  $B$  on  $\bar{X}$  and  $A$  is an exceptional  $\mathbb{R}$ -divisor. We define the *discrepancy*  $a_{K_X+B}(v)$  of  $v$  as the coefficient of  $E$  in  $A - \bar{B}$ . We say that  $(X, B)$  is *log canonical* if  $a_{K_X+B}(v) \geq -1$  for any algebraic valuation  $v$ .

Let  $S$  be a reduced divisor on  $X$  which has no common component with the support of  $B$ , and let  $\nu: S^\nu \rightarrow S$  denote the normalisation of  $S$ . If  $(X, S + B)$  is a log pair, we can construct an effective  $\mathbb{R}$ -divisor  $B^\nu$  on  $S^\nu$ , called the *different* of  $B$  on  $S^\nu$ , in the following manner; see [4, Chapter 16] or [7, §3] for details. The construction is reduced to the case where  $X$  is a surface. Take a resolution of singularities  $f: \bar{X} \rightarrow X$  of the surface  $X$  such that the birational transform  $S^\nu$  of  $S$  on  $\bar{X}$  is also non-singular. Then  $S^\nu$  is the normalisation of  $S$  and we can find an exceptional  $\mathbb{R}$ -divisor  $A$  on  $\bar{X}$  such that  $K_{\bar{X}} + S^\nu + \bar{B} \equiv_f A$ . The different  $B^\nu$  is defined as the  $\mathbb{R}$ -divisor  $(\bar{B} - A)|_{S^\nu}$ . The pair  $(S^\nu, B^\nu)$  is a log pair and it satisfies that  $K_{S^\nu} + B^\nu = \nu^*((K_X + S + B)|_S)$ .

In this paper we prove *inversion of adjunction* on log canonicity as stated below. It was conjectured and proved in dimension three by Shokurov in [7].

**Theorem.** *Let  $(X, S + B)$  be a log pair such that  $S$  is a reduced divisor which has no common component with the support of  $B$ , let  $S^\nu$  denote the normalisation of  $S$ , and let  $B^\nu$  denote the different of  $B$  on  $S^\nu$ . Then  $(X, S + B)$  is log canonical near  $S$  if and only if  $(S^\nu, B^\nu)$  is log canonical.*

Note that this statement has been conjecturally generalised to the precise version of inversion of adjunction in [4, 17.3 Conjecture]. The exposition [5, Section 5.4] is a good reference on inversion of adjunction.

We begin to prove the theorem. The statement is local so we discuss over the germ at a closed point  $x \in S \subset X$ . For a resolution of singularities  $f: \bar{X} \rightarrow X$  which is proper, we call  $f$  a *good resolution* if its exceptional locus on  $\bar{X}$  is a divisor of which the summation with the support of the birational transform of  $S + B$  is a simple normal crossing divisor. Given an ideal sheaf  $\mathfrak{a}$  on  $X$ , we say that  $f$  is a *good resolution for  $\mathfrak{a}$*  if furthermore the ideal sheaf  $\mathfrak{a}\mathcal{O}_{\bar{X}}$  on  $\bar{X}$  is an invertible sheaf. The existence of good resolutions is

guaranteed by Hironaka's resolution theorem [2]. We fix a good resolution of singularities  $f: \bar{X} \rightarrow X$ . Let  $\bar{S} + \bar{B}$  denote the birational transform of  $S + B$  on  $\bar{X}$ ,  $\nu: S^\nu \rightarrow S$  the normalisation morphism, and  $g: \bar{S} \rightarrow S^\nu$  the induced morphism. Let  $\{E_i\}_{i \in I}$  denote the set of prime divisors on  $\bar{X}$  which appears in either the exceptional locus of  $f$  or the support of  $\bar{B}$ , and write  $K_{\bar{X}} + \bar{S} = f^*(K_X + S + B) + \sum_{i \in I} a_i E_i$  with  $a_i := a_{K_X + S + B}(v_{E_i})$ . Then  $K_{\bar{S}} = g^*(K_{S^\nu} + B^\nu) + \sum_{i \in I_S} a_i E_i|_{\bar{S}}$ , where  $I_S := \{i \in I \mid E_i \cap \bar{S} \neq \emptyset\}$ . We mention that  $(X, S + B)$  is log canonical if and only if  $a_i \geq -1$  for any  $i \in I$ , and  $(S^\nu, B^\nu)$  is log canonical if and only if  $a_i \geq -1$  for any  $i \in I_S$ . In particular the only-if-part of the theorem is trivial. It suffices to prove the if-part of the theorem.

From now on we suppose that  $a_i \geq -1$  for any  $i \in I_S$ . Let  $L^\nu$  denote the reduced locus in  $S^\nu$  defined as the union of  $g(E_i|_{\bar{S}})$  for all  $i \in I_S$  with  $a_i = -1$ .  $L^\nu$  is called the *locus of log canonical singularities* for  $(S^\nu, B^\nu)$  and it is independent of the choice of  $f$ . We let  $L$  denote the reduced locus  $\nu(L^\nu)$  and  $\mathfrak{a}_L$  the ideal sheaf on  $X$  defining  $L$ . We construct a sequence of ideal sheaves  $\{\mathfrak{a}_j\}_{j \geq 0}$  on  $X$  in the following manner.

- (i) Set  $\mathfrak{a}_0 := \mathfrak{a}_L$ .
- (ii) Suppose that  $\mathfrak{a}_j$  is given. Take a good resolution  $f_j: X_j \rightarrow X$  for  $\mathfrak{a}_j$  which factors through  $f$ .
- (iii) Let  $S_j + B_j$  denote the birational transform of  $S + B$  on  $X_j$ , and  $\{E_{j,i}\}_{i \in I_j}$  the set of prime divisors on  $X_j$  which appears in either the exceptional locus of  $f_j$  or the support of  $B_j$ . We can regard the set  $I$  of indices as a subset of  $I_j$  by setting  $E_{j,i}$  for  $i \in I$  as the birational transform of  $E_i$ . Set  $I_{j,S} := \{i \in I_j \mid E_{j,i} \cap S_j \neq \emptyset\}$ .
- (iv) Set  $a_i := a_{K_X + S + B}(v_{E_{j,i}})$  for  $i \in I_j$ . Then  $K_{X_j} + S_j = f_j^*(K_X + S + B) + \sum_{i \in I_j} a_i E_{j,i}$ . Write  $\mathfrak{a}_j \mathcal{O}_{X_j} = \mathcal{O}_{X_j}(-\sum_{i \in I_j} n_{j,i} E_{j,i})$ . We set  $\mathfrak{a}_{j+1} := \mathfrak{a}_L \cap f_{j*} \mathcal{O}_{X_j}(\sum_{i \in I_j} (\lfloor a_i \rfloor + 1 - n_{j,i}) E_{j,i})$ . Here  $\lfloor a \rfloor$  denotes the *round-down* of  $a$ , that is, the largest integer which is at most  $a$ .

The below inequality is obvious by the construction of  $\mathfrak{a}_{j+1}$ .

**Lemma 1.**  $n_{j+1,i} \geq n_{j,i} - (\lfloor a_i \rfloor + 1)$  for  $i \in I$ .

Let  $\mathfrak{a}_S$  denote the ideal sheaf on  $X$  defining  $S$ . Kawamata–Viehweg vanishing theorem [3, Theorem 1-2-3] deduces the following equality.

**Lemma 2.**  $\mathfrak{a}_j + \mathfrak{a}_S = \mathfrak{a}_L$  for any  $j$ .

*Proof.* We shall prove the lemma by induction on  $j$ .  $\mathfrak{a}_0 + \mathfrak{a}_S = \mathfrak{a}_L$  is trivial by  $\mathfrak{a}_0 = \mathfrak{a}_L \supset \mathfrak{a}_S$ . Suppose that  $\mathfrak{a}_j + \mathfrak{a}_S = \mathfrak{a}_L$  for fixed  $j$ . Take a small positive rational number  $\epsilon$  so that  $\lceil a_i - (1 - \epsilon)n_{j,i} \rceil = \lfloor a_i \rfloor + 1 - n_{j,i}$  for any  $i \in I_j$  with  $n_{j,i} > 0$ , where  $\lceil a \rceil$  denotes the *round-up* of  $a$ , that is, the smallest integer which is at least  $a$ . We have a natural exact sequence

$$\begin{aligned}
0 \rightarrow \mathcal{O}_{X_j} \left( \sum_{i \in I_j} [a_i - (1 - \epsilon)n_{j,i}] E_{j,i} - S_j \right) \rightarrow \\
\mathcal{O}_{X_j} \left( \sum_{i \in I_j} [a_i - (1 - \epsilon)n_{j,i}] E_{j,i} \right) \rightarrow \mathcal{O}_{S_j} \left( \sum_{i \in I_{j,S}} [a_i - (1 - \epsilon)n_{j,i}] E_{j,i}|_{S_j} \right) \rightarrow 0.
\end{aligned}$$

Since  $\sum_{i \in I_j} (a_i - (1 - \epsilon)n_{j,i})E_{j,i} - S_j \equiv_{f_j} K_{X_j} - (1 - \epsilon)\sum_{i \in I_j} n_{j,i}E_{j,i}$ , we have  $R^1 f_{j*} \mathcal{O}_{X_j}(\sum_{i \in I_j} [a_i - (1 - \epsilon)n_{j,i}]E_{j,i} - S_j) = 0$  by Kawamata–Viehweg vanishing theorem [3, Theorem 1-2-3]. Here we should mention that this theorem [3, Theorem 1-2-3] holds even if  $D$  is an  $\mathbb{R}$ -divisor by its proof. Hence we obtain the surjective map

$$f_{j*} \mathcal{O}_{X_j}(\sum_{i \in I_j} [a_i - (1 - \epsilon)n_{j,i}]E_{j,i}) \rightarrow f_{j*} \mathcal{O}_{S_j}(\sum_{i \in I_{j,S}} [a_i - (1 - \epsilon)n_{j,i}]E_{j,i}|_{S_j}).$$

For any  $i \in I_{j,S}$ , we have  $[a_i - (1 - \epsilon)n_{j,i}] \geq -n_{j,i}$  unless  $a_i = -1$  with  $n_{j,i} = 0$ , and if  $a_i = -1$  then  $f_j(E_{j,i}|_{S_j}) \subset L$ . This implies that

$$\begin{aligned} f_{j*} \mathcal{O}_{S_j}(\sum_{i \in I_{j,S}} [a_i - (1 - \epsilon)n_{j,i}]E_{j,i}|_{S_j}) &\supset f_{j*} \mathcal{O}_{S_j}(-\sum_{i \in I_{j,S}} n_{j,i}E_{j,i}|_{S_j}) \cap \mathbf{a}_L \mathcal{O}_S \\ &\supset \mathbf{a}_j \mathcal{O}_S \cap \mathbf{a}_L \mathcal{O}_S = \mathbf{a}_L \mathcal{O}_S. \end{aligned}$$

Here we used the assumption that  $\mathbf{a}_j + \mathbf{a}_S = \mathbf{a}_L$ . On the other hand  $[a_i - (1 - \epsilon)n_{j,i}] \leq [a_i] + 1 - n_{j,i}$  for any  $i \in I_j$ , whence

$$f_{j*} \mathcal{O}_{X_j}(\sum_{i \in I_j} [a_i - (1 - \epsilon)n_{j,i}]E_{j,i}) \subset f_{j*} \mathcal{O}_{X_j}(\sum_{i \in I_j} ([a_i] + 1 - n_{j,i})E_{j,i}) \subset \mathcal{O}_X.$$

Therefore the above surjective map induces another surjective map

$$\mathbf{a}_{j+1} = \mathbf{a}_L \cap f_{j*} \mathcal{O}_{X_j}(\sum_{i \in I_j} ([a_i] + 1 - n_{j,i})E_{j,i}) \rightarrow \mathbf{a}_L \mathcal{O}_S.$$

Hence  $\mathbf{a}_{j+1} + \mathbf{a}_S = \mathbf{a}_L$  and Lemma 2 is proved. q.e.d.

Let  $\mathfrak{m}$  denote the ideal sheaf on  $X$  defining  $\{x\}$ . The following lemma is due to Zariski's subspace theorem [1, (10.6)].

**Lemma 3.** *Unless  $a_i \geq -1$  for all  $i \in I$ , we can find an integer  $j(l)$  with  $\mathbf{a}_{j(l)} \subset \mathfrak{m}^l$  for each  $l$ .*

*Proof.* We have  $\mathbf{a}_j \subset f_* \mathcal{O}_{\bar{X}}(-n_{j,i}E_i)$  for any  $i \in I$  by the construction of  $\mathbf{a}_j$ . Since  $n_{j,i} \geq n_{0,i} - j([a_i] + 1)$  by Lemma 1, we obtain that  $\mathbf{a}_j \subset f_* \mathcal{O}_{\bar{X}}(-(n_{0,i} - j([a_i] + 1))E_i)$ . Hence Lemma 3 is reduced to finding an integer  $k(l)$  with  $f_* \mathcal{O}_{\bar{X}}(-k(l)E_i) \subset \mathfrak{m}^l$  for fixed  $i$  and  $l$ . Take a closed point  $y$  in  $E_i \cap f^{-1}(x)$ . Then the existence of  $k(l)$  is obtained by Zariski's subspace theorem [1, (10.6)] for  $R := \mathcal{O}_{X,x}$  and  $S := \mathcal{O}_{\bar{X},y}$ . Note that  $\mathcal{O}_{X,x}$  is analytically normal; see [6, (37.5)] for the proof. q.e.d.

We shall complete the proof of the theorem. Unless  $a_i \geq -1$  for all  $i \in I$ , we have  $\bigcap_{j \geq 0} (\mathbf{a}_j + \mathbf{a}_S) = \mathbf{a}_S$  by Lemma 3 and the Artin–Rees lemma; so  $\mathbf{a}_L = \mathbf{a}_S$  by Lemma 2, which is absurd. Therefore  $a_i \geq -1$  for all  $i \in I$  and the theorem is proved.

## REFERENCES

1. S. Abhyankar, *Resolution of singularities of embedded algebraic surfaces*, Second edition, Springer Monographs in Mathematics, Springer-Verlag (1998).
2. H. Hironaka, Resolution of singularities of an algebraic variety over a field of characteristic zero I, II, Ann. of Math. (2) **79** (1964), 109–203; *ibid.* 205–326.
3. Y. Kawamata, K. Matsuda and K. Matsuki, Introduction to the minimal model problem, *Algebraic geometry*, Adv. Stud. Pure Math. **10** (1987), 283–360.

4. J. Kollár et al, *Flips and abundance for algebraic threefolds*, Astérisque **211** (1992).
5. J. Kollár and S. Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Mathematics **134**, Cambridge University Press (1998).
6. M. Nagata, *Local rings*, Interscience Tracts in Pure and Applied Mathematics **13**, Interscience Publishers a division of John Wiley & Sons (1962).
7. V. Shokurov, Three-dimensional log perestroikas, *Izv. Ross. Akad. Nauk Ser. Mat.* **56** (1992), 105-203; translation in *Russian Acad. Sci. Izv. Math.* **40** (1993), 95-202.

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES, KYOTO UNIVERSITY, KYOTO  
606-8502, JAPAN

*E-mail address:* masayuki@kurims.kyoto-u.ac.jp